

LCA Information

Style nr. 19212-291

Contracting organization	Mascot International A/S, Denmark
Project team	Corporate Responsibility Department, Mascot International A/S
Review of Mascot's Life-Cycle Assessment (LCA) methodology and product LCA	Quantis Sàrl, Switzerland
Method validity date	December 2023 Methodology is valid for 5 years
Method	ISO 14040:2006 + A1:2020 / ISO 14044:2006 + A1:2018 + A2:2020. Product Environmental Footprint Category Rules (PEFCR) for Apparel and Footwear is followed when possible.
Description of system boundaries	Cradle to grave
LCIA method	EF 3.1 (adapted)
Data collection	Primary data – main source. Generic data from ecoinvent v.3.10 APOS database Reference year is 2024
LCA software used	SimaPro v.9.6.0.1
Data quality	Method for data quality rating (DQR) developed in alignment with the PEF requirements.
Data quality declaration	High (rated as described in PEFCR for Apparel and Footwear).
Limitations	Style studies are based on reference sizes as defined in PEFCR for apparel and footwear. Current model is also based on reference colours. For other sizes and colours, the reader is encouraged to bear this in mind.
LCA methodology summary report	Contact responsibility@mascot.dk if you are interested in the report.

LIFE CYCLE ASSESSMENT FACTSHEET

April 2025 version 2.1

TARGET GROUP

The 19212 is part of a collection designed for a broad target group in different more demanding work situations within contractors, road workers, offshore and wind sector, airport workers and sailors and with laundry agreements.

LONG-LASTING DURABILITY


By analysing fabric performance requirements and collecting data on customer experience, the LCA is verified by Quantis for an estimated duration of service of use in hard working situations and with industrial wash every week.

CRADLE-TO-GRAVE


Cradle-to-grave is a scoping of the LCA that calculates the entire lifecycle of a product from Extraction of Raw materials to the Use & Wash and End-of-Life stages. Cradle-to-grave results are presented per use according to PEF Category Rules for Apparel and Footwear.

METHODOLOGY

MASCOT LCAs is mainly based on primary data from own factories and suppliers. MASCOT LCAs are calculated according to ISO14040/44. The method is verified by Quantis and applies to all colours.



PROCESS CHAIN



THE 16 IMPACT FACTORS

Impact category	Damage assessment unit	Impact to-gate per garment	Impact to-grave per use
Acidification	mol H ⁺ eq	0,0409	0,000222
Climate change	kg CO ₂ eq	7,71	0,0583
Climate change - Biogenic	kg CO ₂ eq	0,0221	0,000703
Climate change - Fossil	kg CO ₂ eq	7,68	0,0565
Climate change - Land use and LU change	kg CO ₂ eq	0,00961	0,00111
Ecotoxicity, freshwater	CTUe	33,7	0,51
Ecotoxicity, freshwater - part 2	CTUe	27,3	0,143
Ecotoxicity, freshwater - inorganics	CTUe	50,6	0,505
Ecotoxicity, freshwater - organics part 1	CTUe	4,98	0,127
Ecotoxicity, freshwater - organics part 2	CTUe	5,31	0,0215
Particulate matter	disease inc.	0,000000362	0,0000000203
Eutrophication, marine	kg N eq	0,0089	0,0000727
Eutrophication, freshwater	kg P eq	0,00035	0,00000625
Eutrophication, terrestrial	mol N eq	0,0925	0,000545
Human toxicity, cancer	CTUh	0,0000000215	0,00000000177
Human toxicity, cancer - inorganics	CTUh	0,000000000917	0,0000000000562
Human toxicity, cancer - organics	CTUh	0,0000000205	0,000000000171
Human toxicity, non-cancer	CTUh	0,0000000843	0,000000000539
Human toxicity, non-cancer - inorganics	CTUh	0,0000000684	0,000000000476
Human toxicity, non-cancer - organics	CTUh	0,0000000159	0,000000000622
Ionising radiation	kBq U ⁻²³⁵ eq	0,728	0,00262
Land use	Pt	30,5	0,27
Ozone depletion	kg CFC11 eq	0,0000106	0,0000000255
Photochemical ozone formation	kg NMVOC eq	0,0359	0,000207
Resource use, fossils	MJ	125	0,851
Resource use, minerals and metals	kg Sb eq	0,0000973	0,000000358
Water use	m ³ depriv.	6,38	0,024