

LCA Information

Style nr. 19453-316

Contracting organization	Mascot International A/S, Denmark
Project team	Corporate Responsibility Department, Mascot International A/S
Review of Mascot's Life-Cycle Assessment (LCA) methodology and product LCA	Quantis Sàrl, Switzerland
Method validity date	December 2023 Methodology is valid for 5 years
Method	ISO 14040:2006 + A1:2020 / ISO 14044:2006 + A1:2018 + A2:2020. Product Environmental Footprint Category Rules (PEFCR) for Apparel and Footwear is followed when possible.
Description of system boundaries	Cradle to grave
LCIA method	EF 3.1 (adapted)
Data collection	Primary data – main source. Generic data from ecoinvent v.3.10 APOS database Reference year is 2024
LCA software used	SimaPro v.9.6.0.1
Data quality	Method for data quality rating (DQR) developed in alignment with the PEF requirements.
Data quality declaration	High (rated as described in PEFCR for Apparel and Footwear).
Limitations	Style studies are based on reference sizes as defined in PEFCR for apparel and footwear. Current model is also based on reference colours. For other sizes and colours, the reader is encouraged to bear this in mind.
LCA methodology summary report	Contact responsibility@mascot.dk if you are interested in the report.

LIFE CYCLE ASSESSMENT FACTSHEET

April 2025 version 2.1

TARGET GROUP

The 19453 is part of a collection designed for a broad target group in different more demanding work situations within contractors, road workers, offshore and wind sector, airport workers and sailors and with laundry agreements.

LONG-LASTING DURABILITY


By analysing fabric performance requirements and collecting data on customer experience, the LCA is verified by Quantis for an estimated duration of service of use in hard working situations and with industrial wash every week.

CRADLE-TO-GRAVE


Cradle-to-grave is a scoping of the LCA that calculates the entire lifecycle of a product from Extraction of Raw materials to the Use & Wash and End-of-Life stages. Cradle-to-grave results are presented per use according to PEF Category Rules for Apparel and Footwear.

METHODOLOGY

MASCOT LCAs is mainly based on primary data from own factories and suppliers. MASCOT LCAs are calculated according to ISO14040/44. The method is verified by Quantis and applies to all colours.



PROCESS CHAIN



THE 16 IMPACT FACTORS

Impact category	Damage assessment unit	Impact to-gate per garment	Impact to-grave per use
Acidification	mol H ⁺ eq	0,0265	0,000139
Climate change	kg CO ₂ eq	4,91	0,0361
Climate change - Biogenic	kg CO ₂ eq	0,0381	0,000579
Climate change - Fossil	kg CO ₂ eq	4,86	0,0349
Climate change - Land use and LU change	kg CO ₂ eq	0,00673	0,000682
Ecotoxicity, freshwater	CTUe	26,2	0,326
Ecotoxicity, freshwater - part 2	CTUe	17,6	0,0896
Ecotoxicity, freshwater - inorganics	CTUe	38,7	0,328
Ecotoxicity, freshwater - organics part 1	CTUe	3,14	0,0776
Ecotoxicity, freshwater - organics part 2	CTUe	2	0,0101
Particulate matter	disease inc.	0,000000245	0,0000000013
Eutrophication, marine	kg N eq	0,00548	0,0000445
Eutrophication, freshwater	kg P eq	0,00025	0,0000039
Eutrophication, terrestrial	mol N eq	0,0563	0,000332
Human toxicity, cancer	CTUh	0,0000000135	0,000000000109
Human toxicity, cancer - inorganics	CTUh	0,00000000056	0,00000000000347
Human toxicity, cancer - organics	CTUh	0,000000013	0,000000000105
Human toxicity, non-cancer	CTUh	0,0000000431	0,000000000313
Human toxicity, non-cancer - inorganics	CTUh	0,0000000382	0,000000000286
Human toxicity, non-cancer - organics	CTUh	0,00000000491	0,000000000264
Ionising radiation	kBq U ⁻²³⁵ eq	0,573	0,00191
Land use	Pt	22,7	0,175
Ozone depletion	kg CFC11 eq	0,00000583	0,000000014
Photochemical ozone formation	kg NMVOC eq	0,0226	0,000128
Resource use, fossils	MJ	83,1	0,535
Resource use, minerals and metals	kg Sb eq	0,0000652	0,000000231
Water use	m ³ depriv.	4,02	0,015